



Math-Net.Ru

Общероссийский математический портал

A. A. Zamyshlyayeva, S. V. Surovtsev, Численное исследование одной математической модели соболевского типа, *J. Comp. Eng. Math.*, 2015, том 2, выпуск 3, 72–80

DOI: 10.14529/jcem150308

Использование Общероссийского математического портала Math-Net.Ru подразумевает, что вы прочитали и согласны с пользовательским соглашением
<http://www.mathnet.ru/rus/agreement>

Параметры загрузки:

IP: 3.147.79.224

29 сентября 2024 г., 21:09:06



NUMERICAL INVESTIGATION OF ONE SOBOLEV TYPE MATHEMATICAL MODEL

*A.A. Zamyshlyeva*¹, alzama@mail.ru,

*S.V. Survtsev*¹, ssw-92@ya.ru.

¹ South Ural State University, Chelyabinsk, Russian Federation.

The article is devoted to a numerical investigation of the Boussinesq – Løve mathematical model. Algorithm for finding of the numerical solution to the Cauchy – Dirichlet problem for the Boussinesq – Løve equation modeling longitudinal oscillations in a thin elastic rod with regard to transverse inertia was obtained on the basis of a phase space method and by using a finite differences method. This problem can be reduced to the Cauchy problem for the Sobolev type equation of the second order, which is not solvable for arbitrary initial values. The constructed algorithm includes the additional check if initial data belongs to the phase space. The algorithm is implemented as a program in Matlab. The results of numerical experiments are obtained both in regular and degenerate cases. The graphs of obtained solutions are presented in each case.

Keywords: Boussinesq – Løve equation; Cauchy – Dirichlet problem; finite differences method; Sobolev type equation; phase space; conditions of data consistency; system of difference equations; the Thomas algorithm.

Introduction

Consider Boussinesq – Løve equation

$$(\lambda - \Delta)u_{tt} = \alpha(\lambda' - \Delta)u_t + \beta(\lambda'' - \Delta)u, \quad 0 < x < \pi, \quad t > 0 \quad (1)$$

with initial

$$\begin{aligned} u(x, 0) &= \varphi(x), & 0 < x < \pi, \\ u_t(x, 0) &= \psi(x), & 0 < x < \pi, \end{aligned} \quad (2)$$

and boundary

$$u(0, t) = u(\pi, t) = 0, \quad t > 0 \quad (3)$$

conditions. The functions φ, ψ are given, $u = u(x, t)$ is unknown function, $\Delta = \frac{\partial^2}{\partial x^2}$ is one-dimensional Laplace operator. Mathematical model (1) – (3) describes longitudinal oscillations in an elastic rod with inertia [1]. Problem (1) – (3) was studied by Zamyshlyeva A.A. [2] and her students [3–5]. Algorithm of numerical solution of problem (1) – (3) based on modified Galerkin method was constructed in [6].

Mathematical model (1) – (3) can be reduced to the Cauchy problem

$$u(0) = \varphi, \quad \dot{u}(0) = \psi \quad (4)$$

for operator-differential equation

$$A\ddot{u} = B_1\dot{u} + B_0u. \quad (5)$$

If operator A is continuously invertible, then equation (5) is called non-degenerate or regular. Otherwise, in particular when $\ker A \neq \{0\}$, such equation is called a Sobolev

type equation. It is well-known that the Cauchy problem for Sobolev type equation is not solvable with arbitrary initial values. In our opinion, the most productive approach to the study of these equations is the phase space method. Its foundations were laid by G.A. Sviridyuk and T.G. Sukacheva [7] in the study of semilinear Sobolev type equations of the first order.

Concerning (1) – (3) there were considered [6] three cases depending on parameters $\lambda, \lambda', \lambda''$. In the cases when $\lambda \notin \sigma(\Delta)$ and $(\lambda \in \sigma(\Delta)) \wedge (\lambda = \lambda' \neq \lambda'')$ the phase space of equation (1) was constructed and this confirms results [2]. In the case $(\lambda \in \sigma(\Delta)) \wedge (\lambda \neq \lambda')$, that was eliminated in [2], we have the necessary conditions for the unique solvability of (1) – (3) in the form of dependence of functions $\varphi(x)$ and $\psi(x)$, so it is shown that the phase space in the sense of [2] doesn't exist.

1. Approximate Solution of the Problem

In this investigation we present the approximate solution to (1) – (3) that is constructed by using a finite differences method. Construct a net over the whole surface of the rectangle $(0, \pi) \times (0, T)$

$$x_i = h \cdot i, \quad i = \overline{0, N}, \quad h = \frac{\pi}{N}, \quad t_j = \tau \cdot j, \quad j = \overline{0, M}, \quad \tau = \frac{T}{M}$$

and define the grid function

$$u_{i,j} = u(x_i, t_j), \quad i = \overline{0, N}, \quad j = \overline{0, M}.$$

Then

$$u_t(x_i, t_j) = \frac{u_{i,j+1} - u_{i,j-1}}{2\tau}, \tag{6}$$

the second derivative of t :

$$u_{tt}(x_i, t_j) = \frac{u_{i,j+1} - 2u_{i,j} + u_{i,j-1}}{\tau^2}, \tag{7}$$

the second derivative of x :

$$u_{xx}(x_i, t_j) = \frac{u_{i+1,j} - 2u_{i,j} + u_{i-1,j}}{h^2}, \tag{8}$$

$$i = \overline{2, N-1}, \quad j = \overline{2, M-1}.$$

If we substitute (6) – (8) in to (1) and combine like terms, we get:

$$\begin{aligned} & u_{i+1,j+1} \left(-\frac{1}{\tau^2 \cdot h^2} - \frac{\alpha}{2\tau \cdot h^2} \right) + u_{i,j+1} \left(\frac{\lambda}{\tau^2} + \frac{2}{\tau^2 \cdot h^2} + \frac{\alpha\lambda'}{2\tau} + \frac{2\alpha}{2\tau \cdot h^2} \right) + \\ & + u_{i-1,j+1} \left(-\frac{1}{\tau^2 \cdot h^2} - \frac{\alpha}{2\tau \cdot h^2} \right) + u_{i+1,j} \left(\frac{2}{\tau^2 \cdot h^2} - \frac{\beta}{h^2} \right) + \\ & + u_{i,j} \left(-\frac{2\lambda}{\tau^2} - \frac{4}{\tau^2 \cdot h^2} + \beta\lambda'' + \frac{2\beta}{h^2} \right) + u_{i-1,j} \left(\frac{2}{\tau^2 \cdot h^2} - \frac{\beta}{h^2} \right) + \\ & + u_{i+1,j-1} \left(-\frac{1}{\tau^2 \cdot h^2} + \frac{\alpha}{2\tau \cdot h^2} \right) + u_{i,j-1} \left(\frac{\lambda}{\tau^2} + \frac{2}{\tau^2 \cdot h^2} - \frac{\alpha\lambda'}{2\tau} - \frac{2\alpha}{2\tau \cdot h^2} \right) + \\ & + u_{i-1,j-1} \left(-\frac{1}{\tau^2 \cdot h^2} + \frac{\alpha}{2\tau \cdot h^2} \right) = 0, \end{aligned}$$

where $i = \overline{2, N-1}, j = \overline{2, M-1}$. Further, fix j and solve the resultant system of equations for $(j+1)$ -th layer on t by a Thomas algorithm.

2. Algorithm Description

Describe the algorithm in detail. There is one step for every block of algorithm.

Step 1. After the start of program execution it is necessary to input coefficients $\lambda, \lambda', \lambda'', \alpha, \beta$, functions $\varphi(x)$ and $\psi(x)$, length of segment for solvability and time interval $t \in [0, T]$, the number of t and x partitions.

Step 2. Generate searching unknown approximate solution U in loop from 2 to $M - 1$ and in inner loop from 2 to $N - 1$ by the finite differences scheme.

Step 3. Conditional test if λ belongs to Laplace operator spectrum, i.e. if λ can be represented as $-k^2$.

If *step 3* has a true value:

Step 4. Conditional test if $\lambda = \lambda_1$.

If *step 4* has a true value:

Step 5. Conditional test from [6] for initial conditions, i.e. if $(u_0, \varphi_k) = 0$ and $(u_0, \psi_k) = 0$.

Futher, *step 9*.

If *step 4* has a false value:

Step 6. Conditional test from [6] for initial conditions, i.e. if $(u_0, \varphi_k) = (u_0, \psi_k)/\mu_k$, where μ_k is a characteristic equation root for Boussinesq - Ldve equation (1).

Futher, *step 9*.

If *step 5* has a false value:

Step 7. In accordance with condition the program generates a message that there are no solutions.

If *step 6* has a false value:

Step 8. In accordance with condition the program generates a message that there are no solutions.

If *step 5* and *step 6* have a true value and *step 3* has a false value:

Step 9. Generate a system of linear equations in loop from 2 to $M - 1$.

Step 10. Resultant system is solved by a Thomas algorithm in unknown coefficients u_{ij} .

Step 11. Resultant solution is displayed by a graph.

The block diagram of the program is shown on fig 1.

3. Experimental Examples

Example 1. Consider the mathematical model

$$\begin{cases} (\lambda - \Delta)u_{tt} = \alpha(\Delta - \lambda')u_t + \beta(\Delta - \lambda'')u, \\ u(x, 0) = \sin(x), \quad u_t(x, 0) = \sin(x) + \sin(2x), \\ u(0, t) = u(\pi, t) = 0. \end{cases}$$

with parametres $\alpha = -1, \beta = -1, \lambda = 2, \lambda' = -1, \lambda'' = 3$, the amount of steps for x : $N = 40$, for t : $M = 60$. This mathematical model is non-degenerate, therefore the solution exists. The graph of solution is shown on fig. 2.

Example 2. Consider the mathematical model

$$\begin{cases} (\lambda - \Delta)u_{tt} = \alpha(\Delta - \lambda')u_t + \beta(\Delta - \lambda'')u, \\ u(x, 0) = \sin(x), \quad u_t(x, 0) = \sin(x) + \sin(2x), \\ u(0, t) = u(\pi, t) = 0. \end{cases}$$




Fig. 1. The block diagram of the program




Fig. 2. The graph of problem solution from example 1

with parameters $\alpha = -1, \beta = -1, \lambda = -2, \lambda' = -1, \lambda'' = 3$. This mathematical model is non-degenerate, therefore the solution exists. The graph of solution is shown on fig. 3.




Fig. 3. The graph of problem solution from example 2

Example 3. Consider the mathematical model

$$\begin{cases} (\lambda - \Delta)u_{tt} = \alpha(\Delta - \lambda')u_t + \beta(\Delta - \lambda'')u, \\ u(x, 0) = \sin(x), \quad u_t(x, 0) = \sin(x) + \sin(2x), \\ u(0, t) = u(\pi, t) = 0. \end{cases}$$

with parameters $\alpha = -1, \beta = -1, \lambda = -4, \lambda' = -1, \lambda'' = 3$. As $\lambda = \lambda_2$ (it coincides with the second eigenvalue of Laplace operator), the solution doesn't exist because the

conditions of consistency[6]:

$$\frac{1}{\mu_k} \int_0^\pi \psi(x) \cdot \varphi_k(x) dx = \int_0^\pi \varphi(x) \cdot \varphi_k(x) dx$$

doesn't hold. The program gives the message about non-existence of solution shown on fig. 4.




Fig. 4. Message about non-existence of solution in example 3

Example 4. Consider the mathematical model

$$\begin{cases} (\lambda - \Delta)u_{tt} = \alpha(\Delta - \lambda')u_t + \beta(\Delta - \lambda'')u, \\ u(x, 0) = \sin(x), \quad u_t(x, 0) = \sin(x) + \sin(2x), \\ u(0, t) = u(\pi, t) = 0. \end{cases}$$

with parameters $\alpha = -1, \beta = -1, \lambda = -1, \lambda' = -1, \lambda'' = 3$. As $\lambda = \lambda' = \lambda_1$ (coincide with the first eigenvalue of Laplace operator), then for existence of the solution it is necessary that initial functions belong to the phase space of equation, i.e. the following conditions must be satisfied [6]:

$$\int_0^\pi \psi(x) \cdot \varphi_k(x) dx = 0 \text{ and } \int_0^\pi \varphi(x) \cdot \varphi_k(x) dx = 0.$$

Obviously they is don't hold. The program gives the message about non-existence of solution shown on fig. 5.



Fig. 5. Message about non-existence of solution in example 4

Example 5. Consider the mathematical model

$$\begin{cases} (\lambda - \Delta)u_{tt} = \alpha(\Delta - \lambda')u_t + \beta(\Delta - \lambda'')u, \\ u(x, 0) = \sin(3x), \quad u_t(x, 0) = \sin(2x) + \sin(3x), \\ u(0, t) = u(\pi, t) = 0, \end{cases}$$

with parameters $\alpha = -1, \beta = -1, \lambda = -1, \lambda' = -1, \lambda'' = 3$. As $\lambda = \lambda' = \lambda_1$ (coincide with the first eigenvalue of Laplace operator), then for existence of the solution it is necessary that initial functions belong to the phase space of equation, i.e. the following conditions must be satisfied [6]:

$$\int_0^\pi \psi(x) \cdot \varphi_k(x) dx = 0 \text{ and } \int_0^\pi \varphi(x) \cdot \varphi_k(x) dx = 0$$

Obviously they hold. The graph of solution is shown on fig. 6.




Fig. 6. The graph of problem solution from example 5

References

1. Løve A. *A Treatise on the Mathematical Theory of Elasticity*. Moscow, Leningrad, ONTI, 1935. (in Russian)
2. Zamyshlyayeva A.A. The Higher-Order Sobolev-Type Models. *Bulletin of the South Ural State University. Series: Mathematical Modelling, Programming and Computer Software*, 2011, no. 7, p. 5–28. (in Russian)
3. Zamyshlyayeva A.A., Bychkov E.V. The Phase Space of the Modified Boussinesq Equation. *Bulletin of the South Ural State University. Series: Mathematical Modelling, Programming and Computer Software*, 2012, no. 18 (277), issue 12, pp. 13–19. (in Russian)
4. Zamyshlyayeva A.A., Yuzeeva A.V. The Initial-Finish Value Problem for the Boussinesq – Love Equation. *Bulletin of the South Ural State University. Series: Mathematical Modelling, Programming and Computer Software*, 2010, no. 16 (192), issue 5, pp. 23–31. (in Russian)
5. Zamyshlyayeva A.A., Tsyplenkova O.N. The Optimal Control over Solutions of the Initial-finish Value Problem for the Boussinesq – Love Equation. *Bulletin of the South Ural State University. Series: Mathematical Modelling, Programming and Computer Software*, 2012, no. 5 (264), issue 11, pp. 13–24. (in Russian)
6. Zamyshlyayeva A.A., Muravyev A.S. Study of the Mathematical Model Boussinesq–Love. *Vestnik Magnitogorskogo gosudarstvennogo universiteta. Seriya: Matematika* [Bulletin of Magnitogorsk State University. Series: Mathematics], 2013, issue 15, pp. 24–34. (in Russian)
7. Sviridyuk G.A., Sukacheva T.G. [Phase Spaces of One Class of Semilinear Equations of Sobolev Type]. *Differentsial'nyie uravneniia* [Differential Equations], 1990, vol. 26, no. 2, p. 250–258. (in Russian)

Zamyshlyayeva Alyona Aleksandrovna, doctor of physical and mathematical sciences, associate professor, professor of the department of equations of mathematical physics, South Ural State University, Chelyabinsk, Russian Federation, alzama@mail.ru.

Suroutsev Sergey Viktorovich, master student, department of equation of mathematical physics, South Ural State University, Chelyabinsk, Russian Federation, ssw-92@ya.ru.

Received August 20, 2015

УДК 517.9

DOI: 10.14529/jcem150308

ЧИСЛЕННОЕ ИССЛЕДОВАНИЕ ОДНОЙ МАТЕМАТИЧЕСКОЙ МОДЕЛИ СОБОЛЕВСКОГО ТИПА

А.А. Замышляева, С.В. Суворцев

Статья посвящена численному исследованию математической модели Буссинеска – Лява. На основе метода фазового пространства и применения метода конечных разностей построен алгоритм нахождения численного решения задачи Коши – Дирихле для уравнения Буссинеска – Лява, моделирующей продольные колебания в тонком упругом стержне с учетом поперечной инерции. Данная задача может быть редуцирована к задаче Коши для уравнения соболевского типа второго порядка, которая, как известно разрешима не при всех начальных значениях. Разработанный алгоритм содержит предварительную проверку принадлежности начальных данных фазовому пространству. Алгоритм реализован в виде программы в среде Matlab. Приведены результаты вычислительных экспериментов в регулярном и вырожденном случаях. Представлены графики полученных решений.

Ключевые слова: уравнение Буссинеска – Лява; задача Коши – Дирихле; метод конечных разностей; уравнение соболевского типа; фазовое пространство; условия согласования; система разностных уравнений; метод прогонки.

Литература

1. Ляв, А. Математическая теория упругости / А. Ляв. – Москва; Ленинград: ОНТИ, 1935.
2. Замышляева, А.А. Математические модели соболевского типа высокого порядка / А.А. Замышляева // Вестник ЮУрГУ. Серия: Математическое моделирование и программирование. – 2014. – Т. 7, № 2. – С. 5–28.
3. Замышляева, А.А. Фазовое пространство модифицированного уравнения Буссинеска / А.А. Замышляева, Е.В. Бычков // Вестник ЮУрГУ. Серия: Математическое моделирование и программирование. – 2012.– № 18 (277), вып. 12. – С. 13–19.
4. Замышляева, А.А. Начально-конечная задача для уравнения Буссинеска – Лява / А.А. Замышляева, А.В. Юзеева // Вестник ЮУрГУ. Серия: Математическое моделирование и программирование. – 2010. – № 16 (192), вып. 5. – С. 23–31.
5. Замышляева, А.А. Оптимальное управление решениями начально-конечной задачи для уравнения Буссинеска – Лява / А.А. Замышляева, О.Н. Цыпленкова // Вестник ЮУрГУ. Серия: Математическое моделирование и программирование. – 2012.– № 5 (264), вып. 11. – С. 13–24.
6. Замышляева, А.А. Исследование математической модели Буссинеска – Лява / А.А. Замышляева, А.С. Муравьев // Вестник Магнитогорского государственного университета. Серия: Математика. – 2013. – Вып. 15. – С. 24–34.

7. Свиридюк, Г.А. Фазовые пространства одного класса операторных полулинейных уравнений типа Соболева / Г.А. Свиридюк, Т.Г. Сукачева // Дифференциальные уравнения. – 1990.– Т. 26, № 2. – С. 250–258.

Замышляева Алена Александровна, доктор физико-математических наук, доцент, профессор кафедры уравнений математической физики, Южно-Уральский государственный университет (г. Челябинск, Российская Федерация), alzama@mail.ru.

Суровцев Сергей Викторович, магистрант, кафедра уравнений математической физики, Южно-Уральский государственный университет (г. Челябинск, Российская Федерация), ssw-92@ya.ru.

Поступила в редакцию 20 августа 2015 г.