



Math-Net.Ru

Общероссийский математический портал

Waqar Khan, Faisal Yousafzai, Asad Khan, Об упорядоченных группоидах Абеля–Грассмана, *ПФМТ*, 2015, выпуск 2, 40–47

Использование Общероссийского математического портала Math-Net.Ru подразумевает, что вы прочитали и согласны с пользовательским соглашением

<http://www.mathnet.ru/rus/agreement>

Параметры загрузки:

IP: 18.216.149.49

17 ноября 2024 г., 00:06:10



УДК 512.542

ОБ УПОРЯДОЧЕННЫХ ГРУППОИДАХ АБЕЛЯ-ГРАССМАНА

Вагар Хан^{1,2}, Фейсал Ясафзай¹, Асад Хан¹¹Университет науки и технологии Китая, Хэфей, Китай²Институт информационных технологий COMSATS, Абботабад, Пакистан

ON ORDERED ABEL-GRASSMANN'S GROUPOIDS

Waqar Khan^{1,2}, Faisal Yousafzai¹, Asad Khan¹¹University of Science and Technology of China, Hefei, China²COMSATS Institute of Information Technology, Abbottabad, Pakistan

Введено понятие (m, n) -идеалов упорядоченных \mathcal{AG} -группоидов и получены характеристики $(0, 2)$ -идеалов и $(1, 2)$ -идеалов упорядоченного \mathcal{AG} -группоида в терминах левых идеалов. Показано, что упорядоченный \mathcal{AG} -группоид S является $0-(0, 2)$ -бипростым в том и только в том случае, когда S является правым 0 -простым. Результаты данной работы позволяют расширить концепцию \mathcal{AG} -группоида без введенного порядка. Получены характеристики внутренне-регулярного упорядоченного \mathcal{AG} -группоида в терминах левых и правых идеалов.

Ключевые слова: упорядоченные \mathcal{AG} -группоиды, обратимое слева тождество, левая единица, (m, n) -идеал.

The concept of (m, n) -ideals in ordered \mathcal{AG} -groupoids is introduced and the $(0, 2)$ -ideals and $(1, 2)$ -ideals of an ordered \mathcal{AG} -groupoid in terms of left ideals are characterised. It is shown that an ordered \mathcal{AG} -groupoid S is $0-(0, 2)$ -bisimple if and only if S is right 0 -simple. The results of this paper extend the concept of an \mathcal{AG} -groupoid without order. Finally, we characterize an intra-regular ordered \mathcal{AG} -groupoid in terms of left and right ideals.

Keywords: ordered \mathcal{AG} -groupoids, left invertive law, left identity, (m, n) -ideals.

Mathematics Subject Classification (2010): 20D10, 20D20

Introduction

The concept of a left almost semigroup (*LA-semigroup*) [3] was first introduced by M.A. Kazim and M. Naseeruddin in 1972. In [1], the same structure is called a left invertive groupoid. P.V. Protić and N. Stevanović called it an Abel-Grassmann's groupoid (\mathcal{AG} -groupoid) [10].

An \mathcal{AG} -groupoid is a groupoid S satisfying the left invertive law $(ab)c = (cb)a$ for all $a, b, c \in S$. This left invertive law has been obtained by introducing braces on the left of ternary commutative law $abc = cba$. An \mathcal{AG} -groupoid satisfies the medial law $(ab)(cd) = (ac)(bd)$ for all $a, b, c, d \in S$. Since \mathcal{AG} -groupoids satisfy medial law, they belong to the class of entropic groupoids which are also called abelian quasigroups [12]. If an \mathcal{AG} -groupoid S contains a left identity, then it satisfies the paramedial law $(ab)(cd) = (dc)(ba)$ and the identity $a(bc) = b(ac)$ for all $a, b, c, d \in S$ [5].

An \mathcal{AG} -groupoid is a useful algebraic structure, midway between a groupoid and a commutative semigroup. An \mathcal{AG} -groupoid is non-associative and non-commutative in general, however, there is a close relationship with semigroup as well as with

commutative structures. It has been investigated in [5] that if an \mathcal{AG} -groupoid contains a right identity, then it becomes a commutative semigroup. The connection of a commutative inverse semigroup with an \mathcal{AG} -groupoid has been given by Yousafzai et al. in [14] as, a commutative inverse semigroup (S, \cdot) becomes an \mathcal{AG} -groupoid $(S, *)$ under $a*b = ba^{-1}r^{-1}$ for all $a, b, r \in S$. The \mathcal{AG} -groupoid S with left identity becomes a semigroup under the binary operation " \circ_e " defined as, $x \circ_e y = (xe)y$ for all $x, y \in S$ [15]. The \mathcal{AG} -groupoid is the generalization of a semigroup theory [5] and has vast applications in collaboration with semigroups like other branches of mathematics. Many interesting results on \mathcal{AG} -groupoids have been investigated in [7], [8], [9].

If S is an \mathcal{AG} -groupoid with product $\cdot: S \times S \rightarrow S$, then $ab \cdot c$ and $(ab)c$ both denote the product $(a \cdot b) \cdot c$.

Definition 0.1 [16]. An \mathcal{AG} -groupoid (S, \cdot) together with a partial order \leq on S that is compatible with an \mathcal{AG} -groupoid operation, meaning that for $x, y, z \in S$,

$$x \leq y \Rightarrow zx \leq zy \text{ and } xz \leq yz,$$

is called an ordered \mathcal{AG} -groupoid.

Let (S, \cdot, \leq) be an ordered \mathcal{AG} -groupoid. If A and B are nonempty subsets of S , we let

$$AB = \{xy \in S \mid x \in A, y \in B\},$$

and $[A] = \{x \in S \mid x \leq a \text{ for some } a \in A\}$.

Definition 0.2 [16]. Let (S, \cdot, \leq) be an ordered \mathcal{AG} -groupoid. A nonempty subset A of S is called a left (resp. right) ideal of S if the followings hold:

- (i) $SA \subseteq A$ (resp. $AS \subseteq A$);
 - (ii) for $x \in A$ and $y \in S$, $y \leq x$ implies $y \in A$.
- Equivalently $(SA) \subseteq A$ (resp. $(AS) \subseteq A$).

If A is both a left and a right ideal of S , then A is called a two-sided ideal or an ideal of S .

A nonempty subset A of an ordered \mathcal{AG} -groupoid (S, \cdot, \leq) is called \mathcal{AG} -subgroupoid of S if $xy \in A$ for all $x, y \in A$.

It is clear to see that every left and right ideals of an ordered \mathcal{AG} -groupoid is an \mathcal{AG} -subgroupoid.

Let (S, \cdot, \leq) be an ordered \mathcal{AG} -groupoid and let A and B be nonempty subsets of S , then the following was proved in [13]:

- (i) $A \subseteq [A]$;
- (ii) If $A \subseteq B$, then $[A] \subseteq [B]$;
- (iii) $[A][B] \subseteq [AB]$;
- (iv) $[A] = ([A])$;
- (vi) $([A][B]) = [AB]$.

Also for every left (resp. right) ideal T of S , $[T] = T$.

The concept of (m, n) -ideals in ordered semi-groups were given by J. Sanborisoot and T. Changphas in [11]. It's natural to ask whether the concept of (m, n) -ideals in ordered \mathcal{AG} -groupoids is valid or not? The aim of this paper is to deal with (m, n) -ideals in ordered \mathcal{AG} -groupoids. We introduce the concept of (m, n) -ideals in ordered \mathcal{AG} -groupoids as follows:

Definition 0.3. Let (S, \cdot, \leq) be an ordered \mathcal{AG} -groupoid and let m, n be non-negative integers. An \mathcal{AG} -subgroupoid A of S is called an (m, n) -ideal of S if the followings hold:

- (i) $A^m S \cdot A^n \subseteq A$;
- (ii) for $x \in A$ and $y \in S$, $y \leq x$ implies $y \in A$.

Here, A^0 is defined as $A^0 S \cdot A^n = SA^n$ and $A^m S \cdot A^0 = A^m S$.

Equivalently an \mathcal{AG} -subgroupoid A of S is called an (m, n) -ideal of S if

$$(A^m S \cdot A^n) \subseteq A.$$

If $m = n = 1$, then an (m, n) -ideal A of an ordered \mathcal{AG} -groupoid (S, \cdot, \leq) is called a bi-ideal of S .

1 0-minimal $(0, 2)$ -bi-ideals in ordered \mathcal{AG} -groupoid

In this section, we study and generalize the work of W. Jantan and T. Changphas [2] by converting it from an associative ordered structure in to a non-associative ordered structure. We use the concept of (m, n) -ideals and investigate $(0, 2)$ -ideals, $(1, 2)$ -ideals and 0-minimal $(0, 2)$ -ideals in ordered \mathcal{AG} -groupoids. All the results of this section can be obtain for an \mathcal{AG} -groupoid without order.

Definition 1.1. If there is an element 0 of an ordered \mathcal{AG} -groupoid (S, \cdot, \leq) such that $x \cdot 0 = 0 \cdot x = x$ for all $x \in S$, we call 0 a zero element of S .

Example 1.1. Let $S = \{a, b, c, d, e\}$ with a left identity d . Then the following multiplication table and order shows that (S, \cdot, \leq) is a unitary ordered \mathcal{AG} -groupoid with a zero element a .

\cdot	a	b	c	d	e
a	a	a	a	a	a
b	a	e	e	c	e
c	a	e	e	b	e
d	a	b	c	d	e
e	a	e	e	e	e

$$\leq = \{(a, a), (a, b), (c, c), (a, c), (d, d), (a, e), (e, e), (b, b)\}.$$

If S is a unitary ordered \mathcal{AG} -groupoid, then it is easy to see that $(S^2] = S$, $(SA^2] = (A^2S]$ and $A \subseteq (SA] \quad \forall A \subseteq S$. Note that every right ideal of a unitary ordered \mathcal{AG} -groupoid S is a left ideal of S but the converse is not true in general. Example 1.1 shows that there exists a subset $\{a, b, e\}$ of S which is a left ideal of S but not a right ideal of S . It is easy to see that $(SA]$ and $(SA^2]$ are the left and right ideals of a unitary ordered \mathcal{AG} -groupoid S . Thus $(SA^2]$ is an ideal of a unitary ordered \mathcal{AG} -groupoid S .

We characterize of $(0, 2)$ -ideals of an ordered \mathcal{AG} -groupoid in terms of left ideals as follows:

Lemma 1.1. Let (S, \cdot, \leq) be a unitary ordered \mathcal{AG} -groupoid. Then A is a $(0, 2)$ -ideal of S if and only if A is an ideal of some left ideal of S .

Proof. Let A be a $(0, 2)$ -ideal of S , then

$$((SA] \cdot A) = (SA \cdot A) = (AA \cdot S) = (SA^2] \subseteq A,$$

and

$$(A \cdot (SA]) = (A \cdot SA) = (S \cdot AA) = (SA^2] \subseteq A.$$

Hence A is an ideal of a left ideal $(SA]$ of S .

Conversely, assume that A is a left ideal of some left ideal L of S , then

$$(SA^2] = (AA \cdot S) = (SA \cdot A) \subseteq$$

$$\subseteq (SL \cdot A) \subseteq ((SL) \cdot A) \subseteq (LA) \subseteq A,$$

and clearly A is an \mathcal{AG} -subgroupoid of S , therefore A is a $(0, 2)$ -ideal of S .

Corollary 1.1. Let (S, \cdot, \leq) be a unitary ordered \mathcal{AG} -groupoid. Then A is a $(0, 2)$ -ideal of S if and only if A is a left ideal of some left ideal of S .

Now we characterize the $(0, 2)$ -bi-ideals of an ordered \mathcal{AG} -groupoid in terms of right ideals as follows:

Lemma 1.2. Let (S, \cdot, \leq) be a unitary ordered \mathcal{AG} -groupoid. Then A is a $(0, 2)$ -bi-ideal of S if and only if A is an ideal of some right ideal of S .

Proof. Let A be a $(0, 2)$ -bi-ideal of S , then

$$\begin{aligned} ((SA^2] \cdot A] &= (SA^2 \cdot A] = (A^2S \cdot A] = \\ &= (AS \cdot A^2] \subseteq (SA^2] \subseteq A, \end{aligned}$$

and

$$\begin{aligned} (A \cdot (SA^2]) &= (A \cdot SA^2] = \\ &= (A \cdot (S^2]A^2] \subseteq ((A] \cdot (S^2])A^2] \subseteq ((A \cdot S^2A^2]) = \\ &= (A \cdot S^2A^2] = (SS \cdot AA^2] = \\ &= (A^2A \cdot SS] = (SA \cdot A^2] \subseteq (SA^2] \subseteq A. \end{aligned}$$

Hence A is an ideal of some right ideal $(SA^2]$ of S .

Conversely, assume that A is an ideal of some right ideal R of S , then

$$\begin{aligned} (SA^2] &= (A \cdot SA] \subseteq ((A] \cdot (S^2])A] \subseteq \\ &\subseteq ((A \cdot S^2A]) = (A \cdot S^2A] = \\ &= (A \cdot (AS)S] \subseteq (A \cdot (RS)R] \subseteq (A \cdot ((RS)R]) \\ &\subseteq (A \cdot (RS]) \subseteq (AR] \subseteq A, \end{aligned}$$

and $(AS \cdot A] \subseteq ((RS] \cdot A] \subseteq (RA] \subseteq A$, which shows that A is a $(0, 2)$ -ideal of S .

The following result gives some characterizations of $(1, 2)$ -ideals of an ordered \mathcal{AG} -groupoid.

Theorem 1.1. Let (S, \cdot, \leq) be a unitary ordered \mathcal{AG} -groupoid. Then the following statements are equivalent.

- (i) A is a $(1, 2)$ -ideal of S ;
- (ii) A is a left ideal of some bi-ideal of S ;
- (iii) A is a bi-ideal of some ideal of S ;
- (iv) A is a $(0, 2)$ -ideal of some right ideal of S ;
- (v) A is a left ideal of some $(0, 2)$ -ideal of S .

Proof. (i) \Rightarrow (ii): It is easy to see that $(SA^2 \cdot S]$ is a bi-ideal of S . Let A be a $(1, 2)$ -ideal of S , then

$$\begin{aligned} (((SA^2 \cdot S])A] &\subseteq ((SA^2 \cdot SS)A] = \\ &= ((SS \cdot A^2S)A] \subseteq (((S^2] \cdot A^2S)A] = \\ &= ((S \cdot A^2S)A] = ((A^2 \cdot SS)A] \subseteq (A^2S \cdot A] = \\ &= (AS \cdot A^2] \subseteq A, \end{aligned}$$

which shows that A is a left ideal of some bi-ideal $(SA^2 \cdot S]$ of S .

(ii) \Rightarrow (iii): Let A be a left ideal of some bi-ideal B of S and e be a left identity of S , then

$$((A \cdot (SA^2])A] \subseteq ((A \cdot SA^2)A] = ((S \cdot AA^2)A] =$$

$$\begin{aligned} &= e((S \cdot AA^2)A] \subseteq (S)((S \cdot AA^2)A] \subseteq \\ &\subseteq ((S(SA \cdot AA))A] = \\ &= ((S(AA \cdot AS))A] = ((AA \cdot S(AS))A] = \\ &= (((S(AS) \cdot A)A)A] = (((A(SS) \cdot A)A)A] \subseteq \\ &\subseteq (((AS \cdot A)A)A] \subseteq (((BS \cdot B)A)A] \subseteq \\ &\subseteq (BA \cdot A] \subseteq A, \end{aligned}$$

which shows that A is a bi-ideal of an ideal $(SA^2]$ of S .

(iii) \Rightarrow (iv): Let A be a bi-ideal of some ideal I of S , then

$$\begin{aligned} ((SA^2] \cdot A^2] &= (SA^2 \cdot A^2] = ((A^2 \cdot AA)S] = \\ &= ((A \cdot A^2A)S] \subseteq ((A \cdot ((AI)A])S] \subseteq (AA \cdot S] = \\ &= (SA \cdot A] \subseteq ((SI] \cdot S] \subseteq I, \end{aligned}$$

which shows that A is a $(0, 2)$ -ideal of a right ideal $(SA^2]$ of S .

(iv) \Rightarrow (v): It is easy to see that $(SA^3]$ is a $(0, 2)$ -ideal of S . Let A be a $(0, 2)$ -ideal of a right ideal R of S , then

$$\begin{aligned} (A \cdot (SA^3]) &\subseteq (A(SS \cdot A^2A]) \subseteq \\ &\subseteq (A(AA^2 \cdot S]) \subseteq (A((SA \cdot AA)S]) \\ &= (A((AA \cdot AS)S]) = ((AA)((A \cdot AS)S]) \\ &= ((S \cdot A(AS))A^2] = ((A \cdot S(AS))A^2] \\ &\subseteq ((RS] \cdot A^2] \subseteq (RA^2] \subseteq A, \end{aligned}$$

which shows that A is a left ideal of a $(0, 2)$ -ideal $(SA^3]$ of S .

(v) \Rightarrow (i): Let A be a left ideal of a $(0, 2)$ -ideal O of S , then

$$\begin{aligned} (AS \cdot A^2] &\subseteq ((AA \cdot SS)A] \subseteq (SA^2 \cdot A] \subseteq \\ &\subseteq ((SO^2] \cdot A] \subseteq (OA] \subseteq A, \end{aligned}$$

which shows that A is a $(1, 2)$ -ideal of S .

The following characterizes $(1, 2)$ -ideals in terms of left and right ideals of an ordered \mathcal{AG} -groupoid.

Lemma 1.3. Let (S, \cdot, \leq) be a unitary ordered \mathcal{AG} -groupoid and A be an idempotent subset of S . Then A is a $(1, 2)$ -ideal of S if and only if there exist a left ideal L and a right ideal R of S such that $(RL] \subseteq A \subseteq R \cap L$.

Proof. Assume that A is a $(1, 2)$ -ideal of S such that A is idempotent.

Setting $L=(SA]$ and $R=(SA^2]$, then

$$\begin{aligned} (RL] &= ((SA^2] \cdot (SA]) \subseteq (A^2S \cdot SA] \subseteq (A^2S^2 \cdot SA] = \\ &= ((SA \cdot SS)A^2] = \\ &= ((SS \cdot AS)A^2] \subseteq ((S(AA \cdot SS))A^2] = \\ &= ((S(SS \cdot AA))A^2] = \\ &= ((S(A(SS \cdot A)))A^2] \subseteq ((A(S \cdot SA))A^2] \subseteq \\ &\subseteq (AS \cdot A^2] \subseteq A. \end{aligned}$$

It is clear that $A \subseteq R \cap L$.

Conversely, let R be a right ideal and L be a left ideal of S such that $(RL] \subseteq A \subseteq R \cap L$, then

$$(AS \cdot A^2] = (AS \cdot AA] \subseteq ((RS] \cdot (SL]) \subseteq (RL] \subseteq A.$$

Definition 1.2. A $(0,2)$ -ideal A of an ordered \mathcal{AG} -groupoid (S, \cdot, \leq) with zero is said to be 0-minimal if $A \neq \{0\}$ and $\{0\}$ is the only $(0,2)$ -ideal of S properly contained in A .

Remark 1.1. Assume that (S, \cdot, \leq) is a unitary ordered \mathcal{AG} -groupoid with zero. Then it is easy to see that every left (right) ideal of S is a $(0,2)$ -ideal of S . Hence if O is a 0-minimal $(0,2)$ -ideal of S and A is a left (right) ideal of S contained in O , then either $A = \{0\}$ or $A = O$.

Lemma 1.4. Let (S, \cdot, \leq) be a unitary ordered \mathcal{AG} -groupoid with zero. Assume that A is a 0-minimal ideal of S and O is an \mathcal{AG} -subgroupoid of A . Then O is a $(0,2)$ -ideal of S contained in A if and only if $O^2 = \{0\}$ or $O = A$.

Proof. Let O be a $(0,2)$ -ideal of S contained in a 0-minimal ideal A of S . Then $(SO^2] \subseteq O \subseteq A$. Since $(SO^2]$ is an ideal of S , therefore by minimality of A , $(SO^2] = \{0\}$ or $(SO^2] = A$. If $(SO^2] = A$, then $A = (SO^2] \subseteq O$ and therefore $O = A$. Let $(SO^2] = \{0\}$, then

$$(O^2S] \subseteq (O^2S^2] = (S^2O^2] \subseteq (SO^2] = \{0\} \subseteq O^2,$$

which shows that O^2 is a right ideal of S , and hence an ideal of S contained in A , therefore by minimality of A , we have $O^2 = \{0\}$ or $O^2 = A$. Now if $O^2 = A$, then $O = A$.

Conversely, let $O^2 = \{0\}$, then

$$(SO^2] \subseteq (O^2S] = (\{0\}S] = \{0\} = (O).$$

Now if $O = A$, then

$$(SO^2] \subseteq (SS \cdot OO] \subseteq ((SA] \cdot (SA]) \subseteq A = O,$$

which shows that O is a $(0,2)$ -ideal of S contained in A .

Corollary 1.2. Let (S, \cdot, \leq) be a unitary ordered \mathcal{AG} -groupoid with zero. Assume that A is a 0-minimal left ideal of S and O is an \mathcal{AG} -subgroupoid of A . Then O is a $(0,2)$ -ideal of S contained in A if and only if $O^2 = \{0\}$ or $O = A$.

Lemma 1.5. Let (S, \cdot, \leq) be a unitary ordered \mathcal{AG} -groupoid with zero and O be a 0-minimal $(0,2)$ -ideal of S . Then $O^2 = \{0\}$ or O is a 0-minimal right (left) ideal of S .

Proof. Let O be a 0-minimal $(0,2)$ -ideal of S , then

$$(S(O^2)^2] \subseteq (SS \cdot O^2O^2] \subseteq (O^2O^2 \cdot S] = (SO^2 \cdot O^2] \\ \subseteq ((SO^2] \cdot O^2] \subseteq (OO^2] \subseteq O^2,$$

which shows that O^2 is a $(0,2)$ -ideal of S contained in O , therefore by minimality of O , $O^2 = \{0\}$ or $O^2 = O$. Suppose that $O^2 = O$, then

$$(OS] \subseteq (OO \cdot SS] \subseteq (SO^2] \subseteq O,$$

which shows that O is a right ideal of S . Let R be a right ideal of S contained in O , then

$$(R^2S] = (RR \cdot S] \subseteq ((RS] \cdot S] \subseteq R.$$

Thus R is a $(0,2)$ -ideal of S contained in O , and again by minimality of O , $R = \{0\}$ or $R = O$.

The following Corollary follows from Lemma 1.2 and Corollary 1.2.

Corollary 1.3. Let (S, \cdot, \leq) be a unitary ordered \mathcal{AG} -groupoid. Then O is a minimal $(0,2)$ -ideal of S if and only if O is a minimal left ideal of S .

Theorem 1.2. Let (S, \cdot, \leq) be a unitary ordered \mathcal{AG} -groupoid. Then A is a minimal $(2,1)$ -ideal of S if and only if A is a minimal bi-ideal of S .

Proof. Let A be a minimal $(2,1)$ -ideal of S . Then

$$(((A^2S \cdot A])^2S)((A^2S \cdot A)]) \subseteq \\ \subseteq (((A^2S \cdot A)^2S)(A^2S \cdot A)] = \\ = (((A^2S \cdot A)(A^2S \cdot A))S)(A^2S \cdot A)] \subseteq \\ \subseteq (((AS \cdot A)(AS \cdot A))S)(AS \cdot A)] = \\ = (((AS \cdot AS)(AA))S)(AS \cdot A)] \subseteq \\ \subseteq (((A^2S \cdot AA)S)(AS \cdot A)] \subseteq \\ \subseteq (((AS \cdot AS)S)(AS \cdot A)] \subseteq \\ \subseteq ((A^2S \cdot S)(AS \cdot A)] \subseteq \\ \subseteq ((AS \cdot S)(AS \cdot A)] = ((AS \cdot AS)(SA)] \subseteq \\ \subseteq (A^2S \cdot SA] = (AS \cdot SA^2] = ((SA^2 \cdot S)A] \\ \subseteq ((A^2S \cdot S)A] = ((SS \cdot AA)A] = (A^2S \cdot A),$$

and similarly we can show that $(A^2S \cdot A]^2 \subseteq (A^2S \cdot A]$. Thus $(A^2S \cdot A]$ is a $(2,1)$ -ideal of S contained in A , therefore by minimality of A , $(A^2S \cdot A] = A$. Now

$$(AS \cdot A] = ((AS)(A^2S \cdot A)] = \\ = (((A^2S \cdot A)S)A] = ((SA \cdot A^2S)A] = \\ = ((A^2(SA \cdot S))A] \subseteq (A^2S \cdot A] = A,$$

It follows that A is a bi-ideal of S . Suppose that there exists a bi-ideal B of S contained in A , then $(B^2S \cdot B] \subseteq (BS \cdot B] \subseteq B$, so B is a $(2,1)$ -ideal of S contained in A , therefore $B = A$.

Conversely, assume that A is a minimal bi-ideal of S , then it is easy to see that A is a $(2,1)$ -ideal of S . Let C be a $(2,1)$ -ideal of S contained in A , then

$$\begin{aligned} & (((C^2S \cdot C)S)(C^2S \cdot C)] \subseteq \\ & \subseteq (((C^2S \cdot C)S)(C^2S \cdot C)) = \\ & = ((SC \cdot C^2S)(C^2S \cdot C)) = \\ & = ((SC^2 \cdot CS)(C^2S \cdot C)) = \\ & = ((C(SC^2 \cdot S))(C^2S \cdot C)) = \\ & = (((C^2S \cdot C)(SC^2 \cdot SS))C) \subseteq \\ & \subseteq (((C^2S \cdot C)(S \cdot C^2S))C) \subseteq \\ & \subseteq (((C^2S \cdot C)(C^2S))C) \subseteq \\ & = ((C^2((C^2S \cdot C)S))C) \subseteq (C^2S \cdot C). \end{aligned}$$

This shows that $(C^2S \cdot C)$ is a bi-ideal of S , and by minimality of A , $(C^2S \cdot C) = A$. Thus

$$A = (C^2S \cdot C) \subseteq C,$$

and therefore A is a minimal $(2,1)$ -ideal of S .

Theorem 1.3. Let A be 0-minimal $(0,2)$ -bi-ideal of a unitary ordered \mathcal{AG} -groupoid (S, \cdot, \leq) with zero. Then exactly one of the following cases occurs:

- (i) $A = (\{0, a\}]$, $a^2 = 0$;
- (ii) for all $a \in A \setminus \{0\}$, $(Sa^2) = A$.

Proof. Assume that A is a 0-minimal $(0,2)$ -bi-ideal of S . Let $a \in A \setminus \{0\}$, then $(Sa^2) \subseteq A$. Also (Sa^2) is a $(0,2)$ -bi-ideal of S , therefore $(Sa^2) = \{0\}$ or $(Sa^2) = A$.

Let $(Sa^2) = \{0\}$. Since $a^2 \in A$, we have either $a^2 = a$ or $a^2 = 0$ or $a^2 \in A \setminus \{0, a\}$. If $a^2 = a$, then $a^3 = a^2a = a$, which is impossible because $a^3 \in (a^2S) \subseteq (Sa^2) = \{0\}$. Let $a^2 \in A \setminus \{0, a\}$, we have

$$\begin{aligned} & (S \cdot (\{0, a^2\} \{0, a^2\})) \subseteq (SS \cdot a^2a^2) = \\ & = (Sa^2 \cdot Sa^2) = \{0\} \subseteq (\{0, a^2\}], \end{aligned}$$

and

$$\begin{aligned} & (((\{0, a^2\}S)(\{0, a^2\})) \subseteq (\{0, a^2S\} \{0, a^2\}) = \\ & = (a^2S \cdot a^2) \subseteq (Sa^2) = \{0\} \subseteq (\{0, a^2\}]. \end{aligned}$$

Therefore $(\{0, a^2\})$ is a $(0,2)$ -bi-ideal of S contained in A . We observe that $(\{0, a^2\}) \neq \{0\}$ and $(\{0, a^2\}) \neq A$. This is a contradiction to the fact that A is a 0-minimal $(0,2)$ -bi-ideal of S . Therefore $a^2 = 0$ and $A = (\{0, a\}]$. If $(Sa^2) \neq \{0\}$, then $(Sa^2) = A$.

Corollary 1.4. Let A be 0-minimal $(0,2)$ -bi-ideal of a unitary ordered \mathcal{AG} -groupoid (S, \cdot, \leq) with zero such that $(A^2) \neq 0$. Then $A = (Sa^2)$ for every $a \in A \setminus \{0\}$.

Lemma 1.6. Let (S, \cdot, \leq) be a unitary ordered \mathcal{AG} -groupoid. Then every right ideal of S is a $(0,2)$ -bi-ideal of S .

Proof. Assume that A is a right ideal of S , then

$$\begin{aligned} & (Sa^2] \subseteq (AA \cdot SS) \subseteq ((AS) \cdot (AS)) \subseteq \\ & \subseteq (AA) \subseteq (AS) \subseteq A, (AS \cdot A) \subseteq A, \end{aligned}$$

and clearly $A^2 \subseteq A$, therefore A is a $(0,2)$ -bi-ideal of S .

The converse of Lemma 1.2 is not true in general. Example 2.1 shows that there exists a $(0,2)$ -bi-ideal $A = \{a, c, e\}$ of S which is not a right ideal of S .

Definition 1.3. An ordered \mathcal{AG} -groupoid (S, \cdot, \leq) with zero is said to be 0- $(0,2)$ -bisimple if $(S^2) \neq \{0\}$ and $\{0\}$ is the only proper $(0,2)$ -bi-ideal of S .

Theorem 1.4. Let (S, \cdot, \leq) be a unitary ordered \mathcal{AG} -groupoid with zero. Then $(Sa^2) = S$ for all $a \in S \setminus \{0\}$ if and only if S is 0- $(0,2)$ -bisimple if and only if S is right 0-simple.

Proof. Assume that $(Sa^2) = S$ for every $a \in S \setminus \{0\}$. Let A be a $(0,2)$ -bi-ideal of S such that $A \neq \{0\}$. Let $a \in A \setminus \{0\}$, then

$$S = (Sa^2) \subseteq (SA^2) \subseteq A.$$

Therefore $S = A$. Since $S = (Sa^2) \subseteq (S^2)$, we have $(S^2) = S \neq \{0\}$. Thus S is 0- $(0,2)$ -bisimple. The converse statement follows from Corollary 1.2.

Let R be a right ideal of 0- $(0,2)$ -bisimple S . Then by Lemma 1.2, R is a $(0,2)$ -bi-ideal of S and so $R = \{0\}$ or $R = S$. Conversely, assume that S is right 0-simple. Let $a \in S \setminus \{0\}$, then $(Sa^2) = S$. Hence S is 0- $(0,2)$ -bisimple.

Theorem 1.5. Let A be a 0-minimal $(0,2)$ -bi-ideal of a unitary ordered \mathcal{AG} -groupoid (S, \cdot, \leq) with zero. Then either $(A^2) = \{0\}$ or A is right 0-simple.

Proof. Assume that A is 0-minimal $(0,2)$ -bi-ideal of S such that $(A^2) \neq \{0\}$. Then by using Corollary 1.2, $(Sa^2) = A$ for every $a \in A \setminus \{0\}$. Since $a^2 \in A \setminus \{0\}$ for every $a \in A \setminus \{0\}$, we have $a^4 = (a^2)^2 \in A \setminus \{0\}$ for every $a \in A \setminus \{0\}$. Let $a \in A \setminus \{0\}$, then

$$\begin{aligned} & ((Aa^2]S \cdot (Aa^2]) = (a^2A \cdot S(Aa^2)) = \\ & = (((S \cdot Aa^2)A)a^2) \subseteq (((S \cdot A)A)a^2) \\ & \subseteq ((AA \cdot SS)a^2) \subseteq ((SA^2) \cdot a^2) \subseteq (Aa^2], \end{aligned}$$

and

$$\begin{aligned} & (S(Aa^2]^2) = (S((Aa^2] \cdot (Aa^2])) = \\ & = (S((a^2A) \cdot (a^2A))) = (S(a^2(a^2A \cdot A))) = \\ & = ((aa)(S(a^2A \cdot A))) = (((a^2A \cdot A)S)a^2) \subseteq \\ & \subseteq ((AA \cdot SS)a^2) \subseteq ((SA^2) \cdot a^2) \subseteq (Aa^2], \end{aligned}$$

which shows that $(Aa^2]$ is a $(0,2)$ -bi-ideal of S contained in A . Hence $(Aa^2] = \{0\}$ or $(Aa^2] = A$. Since $a^4 \in (Aa^2]$ and $a^4 \in A \setminus \{0\}$, we get $(Aa^2] = A$. Thus by using Theorem 1.2, A is right 0-simple.

2 Ideals in intra-regular ordered \mathcal{AG} -groupoid

Ideal theory plays a very important role in studying and exploring the structural properties of different algebraic structures. Here we study left (right) ideals which usually allow us to characterize an ordered \mathcal{AG} -groupoid and play the role in an ordered \mathcal{AG} -groupoid which is played by normal subgroups in ordered group theory and by ideals in ordered ring theory.

Definition 2.1. An element a of an ordered \mathcal{AG} -groupoid (S, \cdot, \leq) is called an *intra-regular element* of S if there exist some $x, y \in S$ such that $a \leq xa^2 \cdot y$ and S is called *intra-regular* if every element of S is *intra-regular* or equivalently, $A \subseteq (SA^2 \cdot S]$ for all $A \subseteq S$ and $a \in (Sa^2 \cdot S]$ for all $a \in S$.

Example 2.1. Let $S = \{a, b, c, d, e\}$ be an ordered \mathcal{AG} -groupoid with the following multiplication table and order below.

\cdot	a	b	c	d	e
a	a	a	a	a	a
b	a	b	b	b	b
c	a	b	d	e	c
d	a	b	c	d	e
e	a	b	e	c	d

$\leq := \{(a, a), (a, b), (c, c), (d, d), (e, e), (b, b)\}$.

By routine calculation, it is easy to verify that S is intra-regular.

Definition 2.2. An ordered \mathcal{AG} -groupoid (S, \cdot, \leq) is called *left (resp. right) simple* if it has no proper left (resp. right) ideal and is called *simple* if it has no proper ideal.

Theorem 2.1. The following conditions are equivalent for a unitary ordered \mathcal{AG} -groupoid (S, \cdot, \leq) :

- (i) $(aS] = S$, for some $a \in S$;
- (ii) $(Sa] = S$, for some $a \in S$;
- (iii) S is simple;
- (iv) $(AS] = S = (SA]$, where A is any two-sided ideal of S ;
- (v) S is intra-regular.

Proof. (i) \Rightarrow (ii): Let S be a unitary ordered \mathcal{AG} -groupoid and assume that $(aS] = S$ holds for some $a \in S$. Since $(aS]$ and $(Sa]$ are the left ideals of S , then $(aS] = aS$ and $(Sa] = Sa$. Therefore

$$S = (SS] = ((aS] \cdot S] = (aS \cdot S] = (SS \cdot a] = (Sa].$$

(ii) \Rightarrow (iii): Let S be a unitary ordered \mathcal{AG} -groupoid such that $(aS] = S$ holds for some $a \in S$. Suppose that S is not left simple and let L be a proper left ideal of S , then

$$\begin{aligned} (SL] &\subseteq L \subseteq S = \\ &= (SS] \subseteq (Sa \cdot S] \subseteq ((SS \cdot ea)S] = \\ &= ((ae \cdot SS)S] \subseteq ((ae \cdot S)(SS)] = \\ &= ((Se \cdot a)(SS)] = ((SS)(a \cdot Se)] = \\ &= (a(SS \cdot Se)] \subseteq (aS], \end{aligned}$$

implies that $sl \leq at$ for some $a, s, t \in S$ and $l \in L$. Since $sl \in L$, therefore $at \in L$, but $at \in (aS]$. Thus $(aS] \subseteq L$ and therefore we have $S = (aS] \subseteq L$, which implies that $S = L$, which contradicts the given assumption. Thus S is left simple and similarly we can show that S is right simple, which shows that S is simple.

(iii) \Rightarrow (iv): Let S be a simple unitary ordered \mathcal{AG} -groupoid and let A be any two-sided ideal of S , then $A = S$. Therefore, we have $(AS] = (SS] = (SA]$.

(iv) \Rightarrow (v): Let S be a unitary ordered \mathcal{AG} -groupoid such that $(AS] = S = (SA]$ holds for any two-sided ideal A of S . Since $(a^2S]$ is two-sided ideal of S such that $(a^2S \cdot S] = S = (S \cdot a^2S]$. Let $a \in S$, then

$$\begin{aligned} a \in S &= (a^2S \cdot S] \subseteq ((aa \cdot SS)S] = \\ &= ((SS \cdot aa)S] \subseteq (Sa^2 \cdot S], \end{aligned}$$

that is $a \leq (xa^2)y$ for some $x, y \in S$. Thus S is intra-regular

(v) \Rightarrow (i): Let S be a unitary intra-regular ordered \mathcal{AG} -groupoid. Let $a \in S$, then there exist $x, y \in S$ such that $a \leq (xa^2)y$. Thus

$$\begin{aligned} a \leq (xa^2)y &= (ex \cdot aa)y = (aa \cdot ex)y \\ &= (y \cdot ex)(aa) = a((y \cdot ex)a) \in aS, \end{aligned}$$

which shows that $S \subseteq (Sa]$ and $(Sa] \subseteq S$ is obvious. Thus $(Sa] = S$ holds for some $a \in S$.

Corollary 2.1. The following conditions are equivalent for any unitary ordered \mathcal{AG} -groupoid (S, \cdot, \leq) :

- (i) $(aS] = S$, for some $a \in S$;
- (ii) $(Sa] = S$, for some $a \in S$;
- (iii) S is right simple;
- (iv) $(AS] = S = (SA]$, where A is any right ideal of S ;
- (v) S is fully regular.

Corollary 2.2. If (S, \cdot, \leq) is a unitary ordered \mathcal{AG} -groupoid, then the following conditions are equivalent:

(i) $(Sa] = S$, for some $a \in S$;

(ii) $(aS] = S$, for some $a \in S$.

Corollary 2.3. If (S, \cdot, \leq) is a unitary ordered \mathcal{AG} -groupoid, then $(eS] = S = (Se]$ holds for $e \in S$, where e is a left identity of S .

Corollary 2.4. The following conditions are equivalent for any unitary ordered \mathcal{AG} -groupoid (S, \cdot, \leq) :

(i) S is intra-regular;

(ii) $(Sa] = S = (aS]$ for some $a \in S$.

Definition 2.3. A left (resp. right) ideal A of an ordered \mathcal{AG} -groupoid (S, \cdot, \leq) is called semi-prime if $a \in A$ implies $a^2 \in A$.

Lemma 2.1. The following conditions are equivalent for a unitary ordered \mathcal{AG} -groupoid (S, \cdot, \leq) :

(i) S is intra-regular;

(ii) Every right ideal of S is semiprime.

Proof. (i) \Rightarrow (ii): Let T be a right ideal of a unitary intra-regular ordered \mathcal{AG} -groupoid S . For $a \in S$ there exist $x, y \in S$ such that $a \leq xa^2 \cdot y$. Let $a^2 \in T$, then

$$\begin{aligned} a &\leq (ex \cdot a^2)y = (a^2 \cdot xe)y = (y \cdot xe)a^2 = \\ &= a^2(xe \cdot y) \in TS \subseteq (TS) \subseteq T, \end{aligned}$$

which implies that T is semiprime.

Now (ii) \Rightarrow (i): Since $(a^2S]$ is a right ideal of a unitary ordered \mathcal{AG} -groupoid S containing a^2 so $a \in (a^2S]$. Thus

$$\begin{aligned} a \in (a^2S] \subseteq (a^2 \cdot SS] = (S \cdot a^2S] \subseteq (SS \cdot a^2S] = \\ = (Sa^2 \cdot SS] \subseteq (Sa^2 \cdot S]. \end{aligned}$$

Hence S is intra-regular.

Corollary 2.5. The following conditions are equivalent for any unitary ordered \mathcal{AG} -groupoid (S, \cdot, \leq) :

(i) S is intra-regular;

(ii) every ideal of S is semiprime.

Theorem 2.2. The following conditions are equivalent for a unitary ordered \mathcal{AG} -groupoid (S, \cdot, \leq) :

(i) S is intra-regular;

(ii) $L \cap R \subseteq (LR]$ for every semiprime right ideal R and every left ideal L of S ;

(iii) $L \cap R \subseteq (LR \cdot L]$ for every semiprime right ideal R and every left ideal L of S .

Proof. (i) \Rightarrow (iii): Let S be a unitary intra-regular ordered \mathcal{AG} -groupoid and L, R be any left and right ideals of S respectively such that $k \in L \cap R$. Then there exist $x, y \in S$ such that $k \leq xk^2 \cdot y$. Thus

$$\begin{aligned} k &\leq (x \cdot kk)y = (k \cdot xk)y = \\ &= (y \cdot xk)k \leq (y(x(xk^2 \cdot y)))k = \end{aligned}$$

$$\begin{aligned} &= (y(xk^2 \cdot xy))k = (xk^2 \cdot y(xy))k = \\ &= (x(kk) \cdot y(xy))k = \end{aligned}$$

$$\begin{aligned} &= (k(xk) \cdot y(xy))k \in ((R \cdot SL)S)L \subseteq (RL \cdot S)L = \\ &= LS \cdot RL = LR \cdot SL \subseteq LR \cdot L, \end{aligned}$$

which implies that $L \cap R \subseteq (LR \cdot L]$. Also by Lemma 1.3, R is semiprime.

(iii) \Rightarrow (ii): Let R and L be the left and right ideals of S respectively and R be semiprime, then

$$\begin{aligned} L \cap R &= R \cap L \subseteq (RL \cdot R] \subseteq \\ &\subseteq (RL \cdot S] \subseteq (RL \cdot SS] = (SS \cdot LR] \\ &= (L(SS \cdot R))] = (L(RS \cdot S))] \subseteq (L \cdot (RS))] \subseteq (LR]. \end{aligned}$$

(ii) \Rightarrow (i): Since $a \in (Sa]$, which is a left ideal of S , and $a^2 \in (a^2S]$, which is a semiprime right ideal of S , therefore by given assumption $a \in (a^2S]$. Thus

$$\begin{aligned} a \in (Sa] \cap (a^2S] \subseteq ((Sa] \cdot (a^2S))] \subseteq (Sa \cdot a^2S] \subseteq \\ \subseteq (SS \cdot a^2S] = (Sa^2 \cdot SS] \subseteq (Sa^2 \cdot S]. \end{aligned}$$

Hence S is intra-regular.

Lemma 2.2. The following conditions are equivalent for a unitary ordered \mathcal{AG} -groupoid (S, \cdot, \leq) :

(i) S is intra-regular;

(ii) every left ideal of S is idempotent.

Proof. It is simple. We omit the proof.

Theorem 2.3. The following conditions are equivalent for a unitary ordered \mathcal{AG} -groupoid (S, \cdot, \leq) :

(i) S is intra-regular;

(ii) $A = ((SA)^2]$, where A is any left ideal of S .

Proof. (i) \Rightarrow (ii): Let A be a left ideal of a unitary intra-regular ordered \mathcal{AG} -groupoid, then $(SA] \subseteq A$ and by Lemma 1.3, $((SA)^2] = (SA] \subseteq A$. Now $A = (AA] \subseteq (SA] = ((SA)^2]$, which implies that $A = ((SA)^2]$.

(ii) \Rightarrow (i): Let A be a left ideal of S , then $A = ((SA)^2] \subseteq (A^2]$, which implies that A is idempotent and by using Lemma 1.3, S is intra-regular.

REFERENCES

- Holgate, P. Groupoids satisfying a simple invertive law / P. Holgate // Math. Student. – 1992. – Vol. 61. – № 1–4. – P. 101–106.
- Jantan, W. On 0-minimal (0,2)-bi-ideals in ordered semigroups / W. Jantan, T. Changphas // Quasigroups and related Systems. – 2013. – Vol. 21. – P. 83–90.

3. *Kazim, M.A.* On almost semigroups / M.A. Kazim, M. Naseeruddin // Aligarh Bull. Math. – 1972. – Vol. 2. – P. 1–7.
 4. *Lajos, S.* Generalized ideals in semigroups / S. Lajos // Acta Sci. Math. – 1961. – Vol. 22. – P. 217–222.
 5. *Mushtaq, Q.* On LA-semigroups / Q. Mushtaq, S.M. Yusuf // Aligarh Bull. Math. – 1978. – Vol. 8. – P. 65–70.
 6. *Mushtaq, Q.* On locally associative LA-semigroups / Q. Mushtaq, S.M. Yusuf // J. Nat. Sci. Math. – 1979. – Vol. 19. – P. 57–62.
 7. *Mushtaq, Q.* On LA-semigroup defined by a commutative inverse semigroup / Q. Mushtaq, S.M. Yusuf // Mat. Vesnik. – 1988. – Vol. 40. – P. 59–62.
 8. *Mushtaq, Q.* n LA-semigroups with weak associative law / Q. Mushtaq, M.S. Kamran // Scientific Khyber. – 1989. – Vol. 1. – P. 69–71.
 9. *Mushtaq, Q.* Ideals in left almost semigroups / Q. Mushtaq, M. Khan // Proceedings of 4th International Pure Mathematics Conference. – 2003. – P. 65–77.
 10. *Protić, P.V.* AG-test and some general properties of Abel-Grassmann's groupoids / P.V. Protić, N. Stevanović // Pure Mathematics and Applications. – 1995. – Vol. 6. – P. 371–383.
 11. *Sanborisoot, J.* On Characterizations of (m, n) -regular ordered semigroups / J. Sanborisoot, T. Changphas // Far East J. Math. Sci. – 2012. – Vol. 65. – P. 75–86.
 12. *Stevanović, N.* Composition of Abel-Grassmann's 3-bands / N. Stevanović, P. V. Protić // Novi Sad, J. Math. – 2004. – Vol. 34. – P. 175–182.
 13. *Xie, X.Y.* Fuzzy radicals and prime fuzzy ideals of ordered semigroups / X.Y. Xie, J. Tang // Inform. Sci. – 2008. – Vol. 178. – P. 4357–4374.
 14. *Yousafzai, F.* Left regular \mathcal{AG} -groupoids in terms of fuzzy interior ideals / F. Yousafzai, N. Yaqoob, A. Ghareeb // Afrika Matematika. – 2013. – Vol. 24. – P. 577–587.
 15. *Yousafzai, F.* On fully regular \mathcal{AG} -groupoids / F. Yousafzai, A. Khan, B. Davvaz // Afrika Matematika. – 2014. – Vol. 25. – P. 449–459.
 16. *Yousafzai, F.* On fuzzy fully regular ordered \mathcal{AG} -groupoids / F. Yousafzai, A. Khan, V. Amjad, A. Zeb // Journal of Intelligent & Fuzzy Systems. – 2014. – Vol. 26. – P. 2973–2982.
- Research of the first author is supported by the NNSF of China (#11371335).*

Поступила в редакцию 31.10.14.