

2. Appendix

УДК 517.55

On Asymptotic Expansion of the Conormal Symbol of the Singular Operator

Alexander M. Sidorov*

Institute of Mathematics,
Siberian Federal University,
av. Svobodny 79, Krasnoyarsk, 660041
Russia

We study the conormal symbol of the singular Bochner-Martinelli integral on a compact closed surface with conical points S in \mathbb{C}^n and evaluate its asymptotic expansion.

Keywords: singular Bochner-Martinelli operator, conormal symbol, conical point

Introduction

Smooth manifolds with conical points are the simplest singular spaces in the hierarchy of stratified varieties. Differential analysis on such manifolds was perhaps initiated by Kondrat'ev [1] who invented the so-called conormal symbol of a differential operator at a singular point.

1. Integral

Theorem 1 ([4]). *Integral (2) induces a bounded linear operator in $\mathcal{L}^{2,\gamma}(X \times [0, R])$ provided $1 - 2n < \gamma < 0$.*

PROOF. We will consider a smooth hypersurface \mathcal{S} in $\mathbb{C}^n \setminus \{0\}$ with a singular point at the origin given by

$$\mathcal{S} = \{(\varphi(r)x, r) \in \mathbb{R}^{2n} : x \in X, r \in [0, R]\}. \quad (1)$$

Gratefully acknowledge the financial support of the RFBR, grant 05-01-00517, and of the SFU by SM project, grant 45.2007.

References

- [1] V.A.Kondrat'ev, Boundary value problems for elliptic equations in domains with conical points, *Trudy Mosk. Mat. Obshch.* **16**(1967), 209–292.
- [2] A.Sidorov, The Bochner-Martinelli Integral, and Its Applications, Birkhäuser Verlag, Basel et al., 1995.
- [3] N.Tarkhanov, N.Vasilevski, Microlocal Analysis of the Bochner-Martinelli Integral, Preprint 2005/25, Universität Potsdam, 2005, 9 pp.
- [4] S.G.Myslivets, Singular Bochner-Martinelli operator on the hypersurfaces with singularities, *Vestnik of Novosibirsk State Univ.*, (2007), no. 2, 45-57 (Russian).

*e-mail: sidorov@lan.krasu.ru